



Synonym

IL4R,CD124,IL4RA

Source

Mouse IL-4 R alpha, Fc Tag(ILR-M5252) is expressed from human 293 cells (HEK293). It contains AA Ile 26 - Arg 233 (Accession # [NP_001008700](#)).
Predicted N-terminus: Ile 26

Molecular Characterization

IL-4 R alpha(Ile 26 - Arg 233) NP_001008700	Fc(Pro 100 - Lys 330) P01857
--	---------------------------------

This protein carries a human IgG1 Fc tag at the C-terminus.

The protein has a calculated MW of 51 kDa. The protein migrates as 60-66 kDa under reducing (R) condition (SDS-PAGE) due to glycosylation.

Endotoxin

Less than 1.0 EU per µg by the LAL method.

Purity

>95% as determined by SDS-PAGE.

Formulation

Lyophilized from 0.22 µm filtered solution in Tris with Glycine, Arginine and NaCl, pH7.5 with trehalose as protectant.

Contact us for customized product form or formulation.

Reconstitution

Please see Certificate of Analysis for specific instructions.

For best performance, we strongly recommend you to follow the reconstitution protocol provided in the CoA.

Storage


For long term storage, the product should be stored at lyophilized state at -20°C or lower.

Please avoid repeated freeze-thaw cycles.

This product is stable after storage at:

- -20°C to -70°C for 12 months in lyophilized state;
- -70°C for 3 months under sterile conditions after reconstitution.


SDS-PAGE



Mouse IL-4 R alpha, Fc Tag on SDS-PAGE under reducing (R) condition. The gel was stained with Coomassie Blue. The purity of the protein is greater than 95%.

Bioactivity-ELISA

SEC-MALS



The purity of Mouse IL-4 R alpha, Fc Tag (Cat. No. ILR-M5252) is more than 85% and the molecular weight of this protein is around 125-145 kDa verified by SEC-MALS.

[Report](#)

Discounts, Gifts,
and more!




Mouse IL-4 R alpha / CD124 Protein, Fc Tag (MALS verified)

Catalog # ILR-M5252



BIOSYSTEMS
Acro
Surprise Inside!



Immobilized Mouse IL-4, His Tag (Cat. No. IL4-M52H5) at 5 µg/mL (100 µL/well) can bind Mouse IL-4 R alpha, Fc Tag (Cat. No. ILR-M5252) with a linear range of 0.2-16 ng/mL (QC tested).

Background

IL-4 is a pleiotropic cytokine produced by activated Th2 cells and mast cells, and plays a pivotal role in immune responses. The effects of IL-4 are mediated after binding to high affinity receptor complexes present on hematopoietic as well as non-hematopoietic cells. Hematopoietic cellular responses to IL-4 are mediated by a high affinity receptor complex comprised of the 140 kDa IL4R α (CD124) subunit and the 70 kDa common cytokine γ c chain (CD132).

Interleukin 4 Receptor (IL4R) also known as CD124, IL4R α and BSF receptor, is a type I cytokine receptor produced by activated Th2 cells and mast cells, and plays an important role in Th2-biased immune responses, alternative macrophage activation, mucosal immunity, allergic inflammation, tumor progression, and atherogenesis. A soluble form of the encoded IL4R protein can be produced by an alternate splice variant or by proteolysis of the membrane-bound protein, and this soluble form can inhibit IL4-mediated cell proliferation and IL5 upregulation by T-cells. IL4R can alternatively associate with IL-13R α 1 to form the type II receptor which is responsive to both IL4 and IL13. Interleukin-4 receptor has been shown to interact with SHC1.

Clinical and Translational Updates

Discounts, Gifts,
and more!



➤ www.acrobiosystems.com

8/7/2024