



Synonym

MSLN,Mesothelin,MPF

Source

Biotinylated Human Mesothelin (296-580), His,Avitag, premium grade(MSN-H82E9) is expressed from human 293 cells (HEK293). It contains AA Glu 296 - Gly 580 (Accession # [AAH09272.1](#)).

Predicted N-terminus: Glu 296

It is produced under our rigorous quality control system that incorporates a comprehensive set of tests including sterility and endotoxin tests. Product performance is carefully validated and tested for compatibility for cell culture use or any other applications in the early preclinical stage. When ready to transition into later clinical phases, we also offer a custom GMP protein service that tailors to your needs. We will work with you to customize and develop a GMP-grade product in accordance with your requests that also meets the requirements for raw and ancillary materials use in cell manufacturing of cell-based therapies.

Molecular Characterization

Mesothelin(Glu 296 - Gly 580)
AAH09272.1 Poly-his Avi

This protein carries a polyhistidine tag at the C-terminus, followed by an Avi tag (Avitag™).

The protein has a calculated MW of 35.9 kDa. The protein migrates as 40-50 kDa when calibrated against [Star Ribbon Pre-stained Protein Marker](#) under reducing (R) condition (SDS-PAGE) due to glycosylation.

Labeling

Biotinylation of this product is performed using Avitag™ technology. Briefly, the single lysine residue in the Avitag is enzymatically labeled with biotin.

Protein Ratio

Passed as determined by the HABA assay / binding ELISA.

Endotoxin

Less than 0.01 EU per µg by the LAL method.

Sterility

Negative

Mycoplasma

Negative.

Purity

>95% as determined by SDS-PAGE.

Formulation

Lyophilized from 0.22 µm filtered solution in PBS, pH7.4 with trehalose as protectant.

Contact us for customized product form or formulation.

Reconstitution

Please see Certificate of Analysis for specific instructions.

For best performance, we strongly recommend you to follow the reconstitution protocol provided in the CoA.

Storage

For long term storage, the product should be stored at lyophilized state at -20°C or lower.


Please avoid repeated freeze-thaw cycles.

This product is stable after storage at:

- -20°C to -70°C for 12 months in lyophilized state;
- -70°C for 3 months under sterile conditions after reconstitution.


SDS-PAGE

SEC-MALS



Discounts, Gifts,
and more!






The purity of Biotinylated Human Mesothelin (296-580), His,Avitag, premium grade (Cat. No. MSN-H82E9) is more than 85% and the molecular weight of this protein is around 38-53 kDa verified by SEC-MALS.

[Report](#)


Biotinylated Human Mesothelin (296-580), His,Avitag, premium grade on SDS-PAGE under reducing (R) condition. The gel was stained with Coomassie Blue. The purity of the protein is greater than 95% (With [Star Ribbon Pre-stained Protein Marker](#)).

Bioactivity-FACS



2e5 of Anti-MSLN CAR-293 cells were stained with 100µL of 1 µg/mL of Biotinylated Human Mesothelin (296-580), His,Avitag, premium grade (Cat. No. MSN-H82E9) and negative control protein respectively, washed and then followed by PE-SA and analyzed with FACS (QC tested).

Bioactivity-ELISA



Immobilized Anti-Human MSLN MAb at 0.5 µg/mL (100 µL/well) can bind Biotinylated Human Mesothelin (296-580), His,Avitag, premium grade (Cat. No. MSN-H82E9) with a linear range of 0.1-0.78 ng/mL (QC tested).


Discounts, Gifts,
and more!





Evaluation of CAR expression

FACS Analysis of Anti-MSLN CAR Expression



293 cells were transfected with Anti-MSLN-scFv and RFP tag. 2e5 of the cells were first stained with B. Biotinylated Human Mesothelin (296-580), His,Avitag, premium grade (Cat. No. MSN-H82E9, 3 µg/mL) and C. Biotinylated Protein Control, followed by FITC Streptavidin. A. Non-transfected 293 cells and C. Biotinylated Protein Control were used as negative control. RFP was used to evaluate CAR (Anti-MSLN-scFv) expression and FITC was used to evaluate the binding activity of Biotinylated Human Mesothelin (296-580), His,Avitag, premium grade (Cat. No. MSN-H82E9).

Background

Mesothelin (MSLN) is also known as CAK1 antigen, Pre-pro-megakaryocyte-potentiating factor, which belongs to the mesothelin family. Mesothelin / MSLN can be proteolytically cleaved into the following two chains by a furin-like convertase: Megakaryocyte-potentiating factor (MPF) and the cleaved form of mesothelin. Both MPF and the cleaved form of mesothelin are N-glycosylated. Mesothelin / MSLN can interact with MUC16. The membrane-anchored forms of MSLN may play a role in cellular adhesion. MPF potentiates megakaryocyte colony formation in vitro.

Clinical and Translational Updates

Discounts, Gifts,
and more!

