



Synonym

SIGLEC10,MGC126774,PRO940,Siglec10,SLG2

Source

Cynomolgus Siglec-10, His Tag(SI0-C52H5) is expressed from human 293 cells (HEK293). It contains AA Thr 17 - Ala 549 (Accession # [A0A2K5WBX8-1](#)).

Predicted N-terminus: Thr 17

Molecular Characterization

Siglec-10(Thr 17 - Ala 549)
A0A2K5WBX8-1 Poly-his

This protein carries a polyhistidine tag at the C-terminus.

The protein has a calculated MW of 60.2 kDa. The protein migrates as 65-75 kDa under reducing (R) condition (SDS-PAGE) due to glycosylation.

Endotoxin

Less than 1.0 EU per µg by the LAL method.

Purity

>95% as determined by SDS-PAGE.

>90% as determined by SEC-MALS.

Formulation

Lyophilized from 0.22 µm filtered solution in 25 mM MES, 150 mM NaCl, 0.5 M Arginine, pH5.0 with trehalose as protectant.

Contact us for customized product form or formulation.

Reconstitution

Please see Certificate of Analysis for specific instructions.

For best performance, we strongly recommend you to follow the reconstitution protocol provided in the CoA.

Storage


For long term storage, the product should be stored at lyophilized state at -20°C or lower.

Please avoid repeated freeze-thaw cycles.

This product is stable after storage at:

- -20°C to -70°C for 12 months in lyophilized state;
- -70°C for 3 months under sterile conditions after reconstitution.


SDS-PAGE



Cynomolgus Siglec-10, His Tag on SDS-PAGE under reducing (R) condition. The gel was stained with Coomassie Blue. The purity of the protein is greater than 95%.

Bioactivity-ELISA

SEC-MALS



The purity of Cynomolgus Siglec-10, His Tag (Cat. No. SI0-C52H5) is more than 90% and the molecular weight of this protein is around 145-160 kDa verified by SEC-MALS.

[Report](#)

Discounts, Gifts,
and more!




Cynomolgus Siglec-10 Protein, His Tag (MALS verified)

Catalog # SI0-C52H5



BIOSYSTEMS
Acro



Immobilized Cynomolgus Siglec-10, His Tag (Cat. No. SI0-C52H5) at 1 µg/mL (100 µL/well) on a Nickel Coated plate can bind Anti-Siglec-10 Antibody, Human IgG1 with a linear range of 0.009-0.156 µg/mL (QC tested).

Background

The siglecs (sialic acid-binding Ig-like lectins) are a distinct subset of the Ig superfamily with adhesion-molecule-like structure. We describe here a novel member of the siglec protein family that shares a similar structure including five Ig-like domains, a transmembrane domain, and a cytoplasmic tail containing two ITIM-signaling motifs. Siglec-10 was identified through database mining of an asthmatic eosinophil EST library. The Siglec-10-VAP-1 interaction seems to mediate lymphocyte adhesion to endothelium and has the potential to modify the inflammatory microenvironment via the enzymatic end products.

Clinical and Translational Updates

Please contact us via TechSupport@acrobiosystems.com if you have any question on this product.

Discounts, Gifts,
and more!



➤ www.acrobiosystems.com