



Synonym

DIF, TNF-alpha, TNFA, TNFSF2, cachexin, cachectin, TNF α

Source

Human TNF-alpha, premium grade(TNA-H4211) is expressed from human 293 cells (HEK293). It contains AA Val 77 - Leu 233 (Accession # [P01375-1](#)).

Predicted N-terminus: Val 77

It is produced under our rigorous quality control system that incorporates a comprehensive set of tests including sterility and endotoxin tests. Product performance is carefully validated and tested for compatibility for cell culture use or any other applications in the early preclinical stage. When ready to transition into later clinical phases, we also offer a custom GMP protein service that tailors to your needs. We will work with you to customize and develop a GMP-grade product in accordance with your requests that also meets the requirements for raw and ancillary materials use in cell manufacturing of cell-based therapies.

Molecular Characterization

TNF-alpha(Val 77 - Leu 233)
P01375-1

This protein carries no "tag".

The protein has a calculated MW of 17.4 kDa. The protein migrates as 17 kDa \pm 3 kDa when calibrated against [Star Ribbon Pre-stained Protein Marker](#) under reducing (R) condition (SDS-PAGE) due to glycosylation.

Endotoxin

Less than 0.01 EU per μ g by the LAL method.

Host Cell Protein

<0.5 ng/ μ g of protein tested by ELISA.

Host Cell DNA

<0.02 ng/ μ g of protein tested by qPCR.

Sterility

Negative

Mycoplasma

Negative.

Purity

>95% as determined by SDS-PAGE.

>95% as determined by SEC-MALS.

Formulation

Lyophilized from 0.22 μ m filtered solution in PBS, pH7.4 with trehalose as protectant.

Contact us for customized product form or formulation.

Reconstitution

Please see Certificate of Analysis for specific instructions.

For best performance, we strongly recommend you to follow the reconstitution protocol provided in the CoA.

Storage


For long term storage, the product should be stored at lyophilized state at -20 $^{\circ}$ C or lower.

Please avoid repeated freeze-thaw cycles.


This product is stable after storage at:

- -20 $^{\circ}$ C to -70 $^{\circ}$ C for 24 months in lyophilized state;
- -70 $^{\circ}$ C for 3 months under sterile conditions after reconstitution.

SDS-PAGE



SEC-MALS



Discounts, Gifts,
and more!



Human TNF-alpha Protein, premium grade

Catalog # TNA-H4211




BIOSYSTEMS
Acro

Human TNF-alpha, premium grade on SDS-PAGE under reducing (R) and non-reducing (NR) conditions. The gel was stained with Coomassie Blue. The purity of the protein is greater than 95% (With [Star Ribbon Pre-stained Protein Marker](#)).

The purity of Human TNF-alpha, premium grade (Cat. No. TNA-H4211) is more than 95% and the molecular weight of this protein is around 52-62 kDa verified by SEC-MALS. [Report](#)


Bioactivity-Bioactivity CELL BASE

Human TNF-alpha, premium grade induced cytotoxicity in WEHI-13VAR cells




Human TNF-alpha, premium grade (Cat. No. TNA-H4211) induces cytotoxicity effect on the WEHI-13VAR cells in the presence of the metabolic inhibitor actinomycin D. The specific activity of Human TNF-alpha, premium grade is $>2.00 \times 10^7$ IU/mg, which is calibrated against human TNF-alpha WHO International Standard (NIBSC code: 12/154) (QC tested).

Inhibiting effect of Adalimumab in human TNF-alpha, premium grade cytotoxicity assay



Neutralization assay shows that the cytotoxicity effect of human TNF-alpha, premium grade (Cat. No. TNA-H4211) was inhibited by increasing concentration of Adalimumab. The concentration of TNF-alpha used is 20 μ g/mL. The IC50 is 8.48 ng/mL (Routinely tested).

Human TNF-alpha Protein, premium grade stimulates human NF-KB (Luc) HEK293 Reporter Cells



The NF-kB (Luc) HEK293 Reporter Cell was stimulated with serial dilutions of Human TNF-alpha, premium grade (Cat. No. TNA-H4211). The EC50 value of this effect is 0.5143 ng/mL (Routinely tested).

Bioactivity-ELISA

Discounts, Gifts,
and more!





Human TNF-alpha Protein, premium grade

Catalog # TNA-H4211




BIOSYSTEMS
Acro




Immobilized Human TNF-alpha, premium grade (Cat. No. TNA-H4211) at 1 µg/mL (100 µL/well) can bind Humira with a linear range of 0.1-2 ng/mL (QC tested).

Immobilized Human TNF-alpha, premium grade (Cat. No. TNA-H4211) at 2 µg/mL (100 µL/well) can bind Human TNFR1, Fc Tag (Cat. No. TN1-H5251) with a linear range of 0.1-4 ng/mL (Routinely tested).



Immobilized Human TNF-alpha, premium grade (Cat. No. TNA-H4211) at 5 µg/mL (100 µL/well) can bind Human TNFR2, His Tag (Cat. No. TN2-H5227) with a linear range of 0.039-0.313 µg/mL (Routinely tested).

Bioactivity-SPR



Monoclonal Anti-TNF-alpha Antibody, Human IgG1 (13B8) (Cat. No. TNA-AM493) captured on Protein A Chip can bind Human TNF-alpha, premium grade (Cat. No. TNA-H4211) with an affinity constant of 0.267 nM as determined in a SPR assay (Biacore 8K) (Routinely tested).

Discounts, Gifts,
and more!




➤ www.acrobiosystems.com

7/31/2024



Bioactivity-Stability



The Cell based assay shows batch-to-batch consistency between Acro's GMP and PG TNF-alpha.

Background

Tumor necrosis factor alpha (TNF α) is a cytokine produced primarily by monocytes and macrophages. It is found in synovial cells and macrophages in the tissues. The primary role of TNF α is in the regulation of immune cells. TNF α is able to induce apoptotic cell death, to induce inflammation, and to inhibit tumorigenesis and viral replication. Dysregulation of TNF α production has been implicated in a variety of human diseases, including major depression, Alzheimer's disease and cancer. Recombinant TNF α is used as an immunostimulant under the INN tasonermin. TNF α can be produced ectopically in the setting of malignancy and parallels parathyroid hormone both in causing secondary hypercalcemia and in the cancers with which excessive production is associated.

Clinical and Translational Updates

Discounts, Gifts,
and more!

