



Synonym

CD263, DCR1, DCR1-TNFR, LIT, TRAIL-R3, TRAILR3, TRID

Source

Human TRAIL R3 protein, His Tag(TR3-H52H3) is expressed from human 293 cells (HEK293). It contains AA Ala 26 - Pro 235 (Accession # [O14798](#)).

Predicted N-terminus: Ala 26

Molecular Characterization

TRAIL R3(Ala 26 - Pro 235)
O14798 Poly-his

This protein carries a polyhistidine tag at the C-terminus.

The protein has a calculated MW of 24 kDa. The protein migrates as 43-55 kDa when calibrated against [Star Ribbon Pre-stained Protein Marker](#) under reducing (R) condition (SDS-PAGE) due to glycosylation.

Endotoxin

Less than 1.0 EU per µg by the LAL method.

Purity

>95% as determined by SDS-PAGE.

>90% as determined by SEC-MALS.

Formulation

Lyophilized from 0.22 µm filtered solution in PBS, pH7.4 with trehalose as protectant.

Contact us for customized product form or formulation.

Reconstitution

Please see Certificate of Analysis for specific instructions.

For best performance, we strongly recommend you to follow the reconstitution protocol provided in the CoA.

Storage


For long term storage, the product should be stored at lyophilized state at -20°C or lower.

Please avoid repeated freeze-thaw cycles.

This product is stable after storage at:

- -20°C to -70°C for 12 months in lyophilized state;
- -70°C for 3 months under sterile conditions after reconstitution.


SDS-PAGE



Human TRAIL R3 protein, His Tag on SDS-PAGE under reducing (R) condition. The gel was stained with Coomassie Blue. The purity of the protein is greater than 95% (With [Star Ribbon Pre-stained Protein Marker](#)).

Bioactivity-ELISA

SEC-MALS



The purity of Human TRAIL R3 protein, His Tag (Cat. No. TR3-H52H3) is more than 90% and the molecular weight of this protein is around 35-50 kDa verified by SEC-MALS.

[Report](#)

Discounts, Gifts,
and more!




Human TRAIL R3 protein, His Tag (MALS verified)

Catalog # TR3-H52H3



BIOSYSTEMS
Acro



Immobilized Human TRAIL R3 protein, His Tag (Cat. No. TR3-H52H3) at 5 µg/mL (100 µL/well) can bind Human TRAIL, Mouse IgG2a Fc Tag (Cat. No. TRL-H5259) with a linear range of 0.008-0.5 µg/mL (QC tested).

Background

Tumor necrosis factor receptor superfamily member 10C (TNFRSF10C) is also known as Decoy receptor 1 (DcR1), TNF-related apoptosis-inducing ligand receptor 3 (TRAIL receptor 3 or TRAIL-R3), CD263. TNFRSF10C is a decoy receptor for TRAIL (tumor necrosis factor-related apoptosis-inducing ligand), a member of the tumor necrosis factor (TNF) ligand family. This receptor is not capable of inducing apoptosis (lacks a cytoplasmic death domain), and is thought to function as an antagonistic receptor that protects cells from TRAIL-induced apoptosis.

Clinical and Translational Updates

Discounts, Gifts,
and more!



➤ www.acrobiosystems.com

10/28/2024